Provided for non-commercial research and education use. Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

http://www.elsevier.com/copyright

B R A I N R E S E A R C H R E V I E W S 6 3 (2 0 1 0) 4 7 – 5 9



Review

What determines neurogenic competence in glia?

Marcos Romualdo Costa^a, Magdalena Götz^{b,c}, Benedikt Berninger^{b,c,*}

^aEdmond and Lily Safra International Institute of Neuroscience of Natal, 59066-060 Natal, Brazil and Universidade Federal do Rio Grande do Norte, 59072-970 Natal, Brazil

^bInstitute for Stem Cell Research, National Research Center for Environment and Health, Ingolstädter Landstrasse 1, D-85764 Neuherberg, Germany ^cDepartment of Physiological Genomics, Institute of Physiology, Ludwig-Maximilians University Munich, Schillerstrasse 46, D-80336 Munich, Germany

ARTICLE INFO

Article history: Accepted 14 January 2010 Available online 21 January 2010

Keywords: Stem cells Neurogenesis Proneural gene Brain repair Reactive gliosis Müller glia

ABSTRACT

One of the most intriguing discoveries during the last decade of developmental neurobiology is the fact that both in the developing and adult nervous system neural stem cells often turn out to have a glial identity: Radial glia generates neurons in the developing telencephalon of fish, birds and mammals and astro/radial glial stem cells in specialized neurogenic zones give rise to new neurons throughout life. What are the extrinsic signals acting on and the intrinsic signals acting within these glial populations endowing these with a neurogenic potential, whilst most other glia seemingly lack it? Studies on postnatal astroglia shed interesting light on this question as they are the intermediate between neurogenic radial glia and mature parenchymal astrocytes. At least in vitro their decision to acquire a glial fate is not yet irrevocable as forced expression of a single neurogenic transcription factor enables them to transgress their lineage and to give rise to fully functional neurons acquiring specific subtype characteristics. But even bona fide non-neurogenic glia in the adult nervous system can regain some of their radial glial heritage following injury as exemplified by reactive astroglia in the cerebral cortex and Müller glia in the retina. In this review first we will follow the direction of the physiological times' arrow, along which radial glia become transformed on one side into mature astrocytes gradually losing their neurogenic potential, while some of them seem to escape this dire destiny to settle in the few neurogenic oases of the adult brain where they generate neurons and glia throughout life. But we will also see how pathophysiological conditions partially can reverse the arrow of time reactivating the parenchymal astroglia to re-acquire some of the hallmarks of neural stem cells or progenitors. We will close this review with some thoughts on the surprising compatibility of the co-existence of a neural stem cell and glial identity within the very same cell from the perspective of the concept of transcriptional core networks.

© 2010 Elsevier B.V. All rights reserved.

Contents

0165-0173/\$ – see front matter © 2010 Elsevier B.V. All rights reserved.

doi:10.1016/j.brainresrev.2010.01.002

^{*} Corresponding author. Department of Physiological Genomics, Institute of Physiology, Ludwig-Maximilians University Munich, Schillerstrasse 46, D-80336 Munich, Germany. Fax: +49 89 2180 75 216. E-mail address: benedikt.berninger@lrz.uni-muenchen.de (B. Berninger).

3.	Lineag	ge relationships between adult NSC astroglia and embryonic radial glia	49	
4.	The ro	ble of intercellular signalling pathways for the endowment of glia with neurogenic potential	49	
	4.1.	Notch signalling	49	
	4.2.	Sonic hedgehog (Shh) signalling	51	
5.	Postna	atal astroglia: a transition state between neurogenic radial glia and mature parenchymal astrocytes?	51	
	5.1.	Stable reprogramming of postnatal astroglia	52	
	5.2.	Synapse formation by reprogrammed postnatal astroglia	52	
	5.3.	Recruitment of local or SEZ astroglia following injury of the postnatal cortex?	53	
6.	Re-acquisition of a stem or progenitor-like state by mature glia following injury		53	
	6.1.	Reactivation of astroglia following injury of the cerebral cortex	53	
	6.2.	Reactivation of Müller glia following retinal injury	54	
7.	The co	oncept of a transcriptional network underlying neural stemcellness	55	
Ack	Acknowledgment		56	
Refe	References		56	

1. Introduction

The last ten years have seen the rise of specialized glial cells to the rank of neuronal precursors both in the developing and adult nervous system. In the developing telencephalon neuroepithelial cells differentiate into radial glia which in turn generate either directly or indirectly via intermediate progenitors most forebrain neurons (Anthony et al., 2004; Kriegstein and Alvarez-Buylla, 2009; Malatesta et al., 2003; Malatesta et al., 2000; Pinto and Gotz, 2007). While most of the radial glia (RG) eventually transform into parenchymal astrocytes at the end of neurogenesis (Alves et al., 2002; Voigt, 1989), some of them give rise to astroglial stem cells in the adult subependymal zone (SEZ) or ependymal cells (Merkle et al., 2007; Spassky et al., 2005). Both, radial glia and astroglial stem cells display hallmark features of classical astrocytes [for a detailed review on the glial nature of radial glia see (Pinto and Gotz, 2007)]. Thus the question arises as to what endows these glial cells with a neurogenic competence so noticeably lacking in astroglia in most other regions of the brain, such as e.g. the cerebral cortex. What signals are required for the neurogenic endowment of glia and what processes take place in radial glia while they transform into non-neurogenic astrocytes? Does the loss of neurogenic potential occur abruptly or gradually? Is it in fact irreversible? Our studies during the last few years have led to the recognition that astroglial cells at an early stage of postnatal development are not irrevocably fixed in their lineage, but forced expression of single neurogenic transcription factors can render these cells capable of transgressing their own lineage and generating diverse types of neurons, at least in vitro (Berninger et al., 2007a; Heins et al., 2002). However, similar processes might be evoked in the early postnatal brain by damage such as caused by hypoxia (Fagel et al., 2009; Fagel et al., 2006). Finally, partial neurogenic competence can be regained by glial cells in the adult nervous system following injury as they de-differentiate and resume proliferation, as shown in the retina and the cerebral cortex (Buffo et al., 2008; Fischer and Reh, 2001; Karl et al., 2008). Thus it appears that not all the radial glial heritage is spent during early life, but some of it is latently preserved stimulating the hope that it might be possible to unearth this potential for the development of novel strategies for brain repair.

2. Radial glia: the founding fathers of the telencephalon and the adult neurogenic zones

Radial glial cells (RGCs) comprise a specialized cellular population in most regions of the vertebrate brain during restricted developmental periods, the functions of which have been highly disputed since their first description at late 19th century [reviewed in (Rakic, 2003)]. Nowadays, there is a general consensus that RGCs differentiate from neuroepithelial cells acquiring typical astroglial features, as for instance the presence of glycogen storage granules and the expression of astroglial markers, such as e.g. the astrocyte-specific glutamate and aspartate transporter (GLAST), brain lipidbinding protein (BLBP) and tenascin-C [for review see (Pinto and Gotz, 2007)]. In addition, it is now well established that RGCs function both as neuronal and glial progenitors at least in the developing telencephalon (Anthony and Heintz, 2008; Anthony et al., 2004; Malatesta et al., 2003; Malatesta et al., 2000; Miyata et al., 2001; Noctor et al., 2001; Tamamaki et al., 2001). Finally, with completion of most neuro- and gliogenesis they sign responsible for generating ependymal cells as well as particular astro/radial glial cells that function as neural stem cells (NSCs) in the adult brain (Chojnacki et al., 2009; Kriegstein and Alvarez-Buylla, 2009; Merkle et al., 2004). The fact that at least some RGCs are multipotent, i.e. can generate both neuronal and glial progeny, and are capable of selfrenewing cell divisions generating either two new RGCs (symmetric division) or one RGC and a fate-restricted progenitor (asymmetric division) (Miyata et al., 2001; Noctor et al., 2004) indicate that RGCs exhibit defining stem cell hallmarks and are thus often considered as embryonic NSCs (Kriegstein and Alvarez-Buylla, 2009). The generation of adult NSC from RGCs may be considered as a particular case of RGC selfrenewal raising interesting points about the precise lineage relationships between RGCs and adult NSCs. For example, given the evidence that there are virtually no quiescent RGCs during embryonic development (Hartfuss et al., 2001), consequently all adult NSCs must be derived from RGCs that have been previously contributing to neuro- or gliogenesis. Moreover, given the notion that NSCs in the adult SEZ are quite heterogeneous with respect to the distinct progenies they give rise to (Brill et al., 2009; Brill et al., 2008; Hack et al., 2005; Merkle et al., 2007), it can be argued that each class of adult generated neuron and glia may likely descend from a different type of RGC. Indeed there is compelling evidence that the adult SEZ exhibits a similar dorsoventral organization as the developing telencephalon [for review see (Kriegstein and Alvarez-Buylla, 2009)]: for instance, transcription factors involved in interneuron specification in the developing ventral telencephalon such as Dlx2, (Petryniak et al., 2007) are highly expressed in the ventral SEZ (Brill et al., 2008), while transcription factors expressed dorsally such as Pax6 and Neurogenin2 (Neurog2) (Brill et al., 2009; Brill et al., 2008) are well known for their important role in the developing cerebral cortex (Bertrand et al., 2002). This regionalisation of the SEZ has been taken as sign for the construction of the adult SEZ from "building blocks" of different origins (Kohwi et al., 2007), not unlike a mosaic (Merkle et al., 2007; Young et al., 2007). Finally, while the majority of adult NSCs generate neurons, some of them give rise to oligodendrocytes and possibly also non-stem cell astroglia (Hack et al., 2005) leading to two interrelated questions: (i) Do all adult NSCs in vivo generate neuronal and glial progeny, possibly even in a sequential manner, or are we dealing here with distinct sets of neuro- and gliogenic progenitors (which would no longer justify the use of the term "stem cell")? (ii) In the latter case are different adult progenitors derived from different types of RGCs, i.e. adult neurogenic progenitors from neurogenic RGCs and adult gliogenic progenitors from gliogenic RGCs?

3. Lineage relationships between adult NSC astroglia and embryonic radial glia

Astro/radial glial cells functioning as NSCs in the adult brain have been long suggested to derive from embryonic radial glia (Alvarez-Buylla et al., 2001; Chojnacki et al., 2009). However, embryonic RGCs are heterogeneous (Malatesta et al., 2000) and it is not clear whether adult NSCs would belong to the lineage of multipotent (Fig. 1, scenario I), lineage-restricted (Fig. 1, scenarios II and III) or even of a very small and hence previously missed set of quiescent RGCs (Fig. 1, scenario IV). Recently, we have provided additional evidence indicating that the early developing cerebral cortex is devoid of gliarestricted progenitors (Costa et al., 2009). By using retrovirallybased transduction and video time-lapse microscopy to follow the progeny of single cortical progenitors in vitro and in vivo, we showed that the majority of early cortical progenitors generate exclusively neurons, a pool which becomes depleted with time, whereas a subset of cortical progenitors generate neurons before giving rise to glia-restricted progenitors observed after mid-neurogenesis. Therefore, every actively dividing cortical progenitor seems to generate neurons at early developmental stages and only a subset of progenitors undergoes the alleged sequence of transitions from neurogenesis to gliogenesis during cortical development (Qian et al., 2000). As RGCs have been shown to constitute the majority of progenitor cells in the developing cerebral cortex at the developmental stages when most of these lineage studies were performed (Hartfuss et al., 2001; Malatesta et al., 2000; Noctor et al., 2002, 2004), most of the cortical progenitors studied in previous work [see (Costa et al., 2009) and references herein] are RGCs and, therefore, cell lineage data would indicate that early RGCs comprise two major populations, namely a neuron-restricted and a multipotent population. Since the former seem to become depleted over time as they generate neurons, this may suggest that adult NSCs are encompassed in the lineage of multipotent RGCs contributing both neurons and macroglial cells during development (Alvarez-Buylla et al., 2001) (Fig. 1, scenario I). Although this may be considered to be the most likely scenario, there is no direct experimental support to dismiss the possibility that adult NSCs are derived from symmetrically dividing RGCs generating only daughter RGCs (Fig. 1, scenario III) or from a small, and therefore difficult to be detected, subpopulation of RGCs that did not divide at all during development (Fig. 1, scenario I). None of these alternatives has been directly tested, yet the fact that several signalling pathways involved in the transition from neurogenesis to gliogenesis in the developing cerebral cortex also play a role in the specification of adult NSCs (see below) may lend some support to the view that adult NSCs are indeed encompassed in the lineage of multipotent radial glia (Fig. 1, scenario I).

4. The role of intercellular signalling pathways for the endowment of glia with neurogenic potential

Irrespective of the exact lineage relationship between RGCs and adult NSCs, there is good experimental support for the notion that embryonic RGCs transform into adult SEZ astroglia at postnatal stages (Merkle et al., 2007, 2004). These two cell populations share an intriguing property: they exhibit astroglial features and at the same time have the potential to generate neurons. In contrast, other astroglial population directly derived from RGCs, such as parenchymal astroglia in the cerebral cortex (Alves et al., 2002; Voigt, 1989) and ependymal cells (Spassky et al., 2005) do not generate neurons in the adult brain under physiological conditions. What are the factors endowing RGCs and SEZ astro/radial glia with neurogenic potential and what restricts parenchymal astroglia and ependymal cells to their lineage and prevents them from generating neurons notwithstanding their close ties to RGCs and SEZ astro/radial glia? To gain some more insights into the differences between stem cell and parenchymal glia we need to have a closer look at the signalling pathways activated in these cells. We will focus here on the discussion of Notch and sonic hedgehog (Shh) signalling as several lines of evidence suggest that these pathways play a major role in stem cell fate determination and may hence shed light on the question how astroglial cells are endowed with stem cell properties.

4.1. Notch signalling

On first sight, the canonical Notch pathway seems to be gliogenic: Forced expression of Hes1, a downstream target and effector of Notch, impairs neurogenesis in the embryonic cortex (Ishibashi et al., 1994), whereas expression of an active form of Notch1 (NICD1) or Notch 3 (NICD3) promotes increased generation of parenchymal astrocytes and ependymal cells (Dang et al., 2006; Gaiano et al., 2000). These effects of Notch

Author's personal copy

B R A I N R E S E A R C H R E V I E W S 6 3 (2010) 47-59



Fig. 1 – Possible lineage relationships between RGCs and adult NSCs. In the first scenario (I) RGCs generate neurons and macroglial cells during development before becoming adult NSCs. In the second scenario (II), RGCs that give rise to adult NSCs generate only neurons during development. The third scenario (III) suggests that adult NSCs are derived from a separate population of RGCs that do not contribute neurons or macroglial cells during development. The two lineage trees depicted in each of these 3 scenarios are fictious and have the purpose to illustrate that RGCs may progress to adult NSCs via symmetric and asymmetric cell divisions. The last scenario (IV) suggests the existence of quiescent RGCs that do not divide during development and directly become adult NSCs.

signalling in RGCs have been thought to depend on the repression of proneural genes such as Neurogenin1 (Neurog1), Neurog2 and the mouse homologue of achaete–scute 1(Mash1) by the Notch effectors Hes1 and Hes5 (Hatakeyama et al., 2004). However, it has also been shown that most RGCs express Notch target genes at early embryogenesis (Mizutani et al., 2007), indicating that Notch signalling occurs at stages when RGCs are actively engaged in the generation of neuronal progeny. Moreover, expression of NICD1 or NICD3 significantly increases the frequency of embryonic telencephalic progenitors that form neurospheres (Dang et al., 2006; Yoon et al., 2004), suggesting that Notch signalling might be involved in the maintenance of a stem cell state in RGCs. Accordingly, active Notch 1 and 3 promote the generation of subependymal

astrocytes (Dang et al., 2006; Yoon et al., 2004). Although these data seem at first glance contradictory with previous findings pointing to a gliogenic role for Notch, it might only highlight a limitation inherent to studies based on forced expression or deletion of genes, namely their "all or none" effect that is blind to the temporal fine tuning of physiological expression levels. In fact, it has been recently demonstrated that the levels of Hes1 and Neurog2 oscillate out of phase to each other within RGCs (Shimojo et al., 2008), indicating that the susceptibility of RGCs for fate decisions is dynamically regulated and may thus depend on temporally coincident signalling events impinging on them rather than on the static activation of isolated signalling pathways, such as mimicked by forced expression of Notch effectors or their counter-players. Indeed, promotion of gliogenesis by Notch depends on the concomitant activation of the janus kinase (JAK)-signal transducer and activator of transcription (STAT) pathway (Ge et al., 2002). When these two pathways are not simultaneously activated, the Notch signalling mediator CBF1 binds to a repressive cofactor protein, NCoR, which functions to repress gliogenic genes (Hermanson et al., 2002). Therefore, Notch signalling in embryonic RGCs may prevent neurogenesis by two different means: i) by maintaining RGCs in an undifferentiated, proliferative state and ii) by promoting gliogenesis. In the adult SEZ, Notch signalling functions to maintain ependymal cells in a quiescent state and inhibition of this signalling allows these cells to produce olfactory bulb neurons (Carlen et al., 2009). In contrast, activation of Notch signalling has also been shown to be important for survival and proliferation of adult NSCs (Alexson et al., 2006; Androutsellis-Theotokis et al., 2006), suggesting that similarly to the embryonic brain, the role of Notch signalling in the adult neurogenic niches may also depend on dynamic interactions with other signalling pathways. Interestingly, Notch signalling becomes activated in the cerebral cortex after stroke (Arumugam et al., 2006), a condition in which parenchymal astroglia de-differentiate and resume proliferation and can generate multipotent neurospheres (Buffo et al., 2008, 2005) (see discussion below).

4.2. Sonic hedgehog (Shh) signalling

The Shh member of the Hedgehog protein family plays critical roles in the patterning of the developing neural tube and in the induction of ventral forebrain structures [for a recent review see: (Dessaud et al., 2008)]. Recently, it has been shown that conditional knockout of the Shh and Smo genes in the early developing dorsal telencephalon resulted in a smaller dorsal telencephalon by prolonging the cell cycle of RGCs (Komada et al., 2008). Moreover, Shh signalling has been shown to be crucial for the generation of NSCs in the postnatal brain (Palma et al., 2005). Consistent with a role in neural stem cell maintenance, Shh signalling in the adult neurogenic regions is important to maintain normal levels of NSC proliferation and neurogenesis (Balordi and Fishell, 2007; Han et al., 2008; Lai et al., 2003). Furthermore, in an elegant study, Ahn and Joyner provided unequivocal evidence that a small population of cells (probably RGCs) in the developing brain respond to Shh and establish the two major neurogenic compartments in the adult brain (Ahn and Joyner, 2005). Using a genetic fatemapping strategy based on the expression of the Shh target Gli1 in vivo, they showed that Shh responding cells are established at late embryogenesis and comprise NSCs capable of self-renewal and generation of multiple cell types in the adult brain. More recently, it has also been shown that NSCs fail to develop in animals with defective Shh signalling caused by embryonic ablation of ciliary genes in RGCs (Han et al., 2008). Collectively, these data suggest a pivotal role of Shh for the establishment of NSCs within the neurogenic germinative zones of the adult brain. Surprisingly, yet consistent with such role, Shh can stimulate differentiated parenchymal astroglia from the adult cerebral cortex to initiate the formation of multipotent neurospheres in vitro (Jiao and Chen, 2008). The latter data suggest that this factor may not only be important for the establishment and maintenance of astroglial NSCs

within the stem cell niche, but rather may exert a stem cell identity inducing activity over other astroglial cell populations.

Other molecular determinants may also contribute to the neurogenic potential of stem cell astroglia. For example, members of the bone morphogenetic protein family (BMPs) promote neuronal differentiation at early embryogenesis (Li et al., 1998) and BMP signalling is also active in GFAP-positive cells in the adult SEZ where it is required for neurogenesis (Colak et al., 2008). Likewise, Wnt signalling has also been shown to be involved in the control of cortical progenitors/ RGCs proliferation (Chenn and Walsh, 2002; Woodhead et al., 2006) and adult neurogenesis (Kuwabara et al., 2009; Lie et al., 2005). The precise role of these factors in maintaining the pool of NSCs in developing and in the adult neurogenic germinative zones is currently under active investigation and lessons learned from these studies may help to reactivate neurogenic programs in classically non-neurogenic areas of the brain, as illustrated by the case of Shh.

5. Postnatal astroglia: a transition state between neurogenic radial glia and mature parenchymal astrocytes?

As previously discussed, while some RGCs give rise to astroglial stem cells of the adult neurogenic zones, ependymal cells or late oligodendroglial progenitor cells (Kessaris et al., 2006), the majority is thought to transform directly or indirectly into parenchymal astroglia (Kriegstein and Alvarez-Buylla, 2009). Eventually many of these astroglial cells will take up functions quite different from generating neurons such as regulation of blood flow, ion homeostasis, energy metabolism and regulation of synaptic function (for review see articles within this issue of Brain Research Reviews), i.e. intricate functions which require a high degree of specialization (Wang and Bordey, 2008). Very little is known about the molecular mechanisms underlying the specialization of astroglia, but evidence from the spinal cord white matter points to the possibility that transcription factors involved in neuronal specification also contribute to astrocyte specification, with Pax6 being a prime example for a transcription factor not only regulating neuronal, but also astroglial specification (Hochstim et al., 2008). For instance, astroglia in the ventral and dorsal spinal cord white matter can be distinguished by their mutually exclusive expression of reelin and Slit1, respectively. Intriguingly, reelin requires expression of Pax6, while Nkx 6.1 promotes expression of Slit1 (Hochstim et al., 2008). This data suggest a transcriptional code for astrocyte positional identity in the white matter of the spinal cord. Does expression of neurogenic fate determinants play a similar role in the specification of cortical astroglia? Indeed Pax6 has been detected in postnatal astrocytes and their progenitors (Sakurai and Osumi, 2008) (S. Gascón and M. Götz, unpublished observation) suggesting that continued expression of neurogenic fate determinants - most likely at drastically reduced levels - by cortical astrocytes at early postnatal stages ensures a gradual change from neurogenic RGCs to gliogenic RGCs and finally astrocytes. Indeed several lines of evidence point to such a gradual change: for instance, postnatal astroglia can be induced to initiate the formation of B R A I N R E S E A R C H R E V I E W S 6 3 (2010) 47-59

clonal aggregates called neurospheres when cultured in the presence of epidermal growth factor (EGF) and fibroblast growth factor 2(FGF2) (Laywell et al., 2000) (Heinrich et al., unpublished observation). These neurospheres are characterized by their ability to self-renew (i.e. can give rise to secondary, tertiary etc. neurospheres) and are multipotent, i.e. can generate all three major neural lineages, thus exhibiting the hallmarks of neurospheres generated by NSCs. The neurosphere-forming capacity of astroglia rapidly declines during the second postnatal week (Laywell et al., 2000). Thus, under the influence of EGF/FGF2 early postnatal astroglia can sufficiently dedifferentiate to revert to a multipotent lineage. But by postnatal days 14-18 apparently changes take place within astroglia that disable the neurosphere-forming capacity of these cells. Conceptually, one may assume that these changes correspond with the in parallel increasing involvement of astroglia in their above mentioned regulatory functions. For instance, by about 2 weeks astroglia acquire a mature expression of inwardly rectifying potassium channels (Kir), thought to be involved in buffering of potassium (Bordey and Sontheimer, 1997; Seifert et al., 2009). Interestingly, up-regulation of K_{ir}4.1 has been associated with cell cycle exit, while loss of functional K_{ir} channels is associated with re-entry into cell cycle and gliosis (Olsen and Sontheimer, 2008). Indeed, functional specialization of astroglia is likely to require these cells to cease proliferating and to acquire a postmitotic status thereby rendering these cells refractory to the mitogenic action of EGF and FGF2. However, as we will see below, astroglia can de-differentiate and resume proliferation following injury, an effect which is accompanied by the regaining of neurosphere-forming capacity (Buffo et al., 2008, 2005).

5.1. Stable reprogramming of postnatal astroglia

Another striking example for the plasticity of early postnatal astroglia can be seen in their high susceptibility for neuronal reprogramming following forced expression of neurogenic fate determinants. Based on the fact that neurogenic RGCs in the developing cortex express high levels of Pax6 (Gotz et al., 1998), our laboratory has demonstrated that forced reexpression of Pax6 in culture via retroviral vectors can drive some mouse astroglia towards the neuronal lineage (Heins et al., 2002). Subsequently, we found that retrovirally-mediated expression of the Pax6 target gene Neurog2 or the related proneural gene Mash1 are even more efficient in driving early cultured postnatal astroglia towards neurogenesis (Berninger et al., 2007a). Intriguingly, forced expression of these transcription factors does not only induce expression of neuronal markers, but also results in the gradual acquisition of neuronal conductances and the ability to generate repetitive action potentials discharges suggesting that forced expression of these transcription factors had caused early postnatal astroglia to undergo a stable lineage transgression.

Single cell tracking of astroglia derived from transgenic mice expressing GFP under the human GFAP promoter revealed interesting details about the reprogramming process. First of all, consistent with the viral integration, cells undergoing reprogramming typically divided before undergoing neuronal metamorphosis with many cells apparently dying from apoptosis (Berninger et al., 2007a). It is currently not clear whether this apoptosis is a specific mechanism for aborting failed reprogramming, but it is conceivable that forced expression of neurogenic fate determinants may induce a transcriptional program that enters in catastrophic conflict with the intrinsic astroglial differentiation. Another lesson learned from these single cell tracking experiments is the fact that cells undergoing successful reprogramming exhibit morphological changes surprisingly similar to early cortical progenitors. For instance, like early cortical precursors (LoTurco and Bai, 2006), astroglial cells expressing Neurog2 seem to pass through distinct morphological stages, such as a multipolar followed by a bipolar phase (Fig. 2), and eventually enter a migratory stage (Berninger et al., 2007a). These data are consistent with the recent finding that one of Neurog2's direct transcriptional targets is the small GTP binding protein Rnd2, the activity of which is constitutive and plays a fundamental role in regulating the morphology and migration of early cortical precursors (Heng et al., 2008).

5.2. Synapse formation by reprogrammed postnatal astroglia

Notably, astroglia-derived neurons were capable of receiving functional synapses from co-cultured embryonic cortical neurons suggesting a remarkable ability for functional integration. Somewhat surprisingly, however, astroglia-derived neurons failed to form functional presynaptic outputs. Yet, consistent with Neurog2 playing a fundamental role in specifying a glutamatergic identity in telencephalic precursors (Bertrand et al., 2002), a substantial proportion of astroglia expressing Neurog2, but not Mash1 or Pax6, upregulated the T-box transcription factor Tbr1 (Berninger et al., 2007a), a hallmark in glutamatergic neurogenesis (Hevner et al., 2006). However, the failure of forming functional presynaptic specializations argued that neuronal reprogramming remained only partial. In an effort to overcome these limitations by using viral vectors providing higher levels of expression and less susceptible to silencing we have been recently able to show that early postnatal astroglia can be directed towards fully functional synapse forming neurons, at least in vitro (Heinrich et al., unpublished observation). Importantly, this study showed that forced expression of distinct fate determinants such as Neurog2 and Dlx2, known for their critical roles in specifying glutamatergic and GABAergic neurons, respectively, within the developing telencephalon (Bertrand et al., 2002; Petryniak et al., 2007) directs astroglial cells towards distinct neuronal transmitter subtypes (Heinrich et al., unpublished observation), consistent with their developmental role. Thus, forced expression of neurogenic fate determinants can result in the stable and complete lineage reprogramming of early postnatal astroglia. Notably, in parallel to the decline in neurosphere-forming capacity, the susceptibility of undergoing reprogramming gradually decreases with the age of the cultured astroglia (Berninger, unpublished observation) again arguing for a gradual change in the internal milieu of astroglia undergoing maturation. Future studies will have to show whether these changes are accounted for by epigenetic modifications and whether and under what conditions they are reversible.



Fig. 2 – Metamorphosis of an astrocyte from the postnatal cerebral cortex into neuron upon forced expression of Neurog2. Panels depict different stages of cellular siblings (red arrows) derived from a hGFAP-GFP positive postnatal astroglia transduced with Neurog2. Time is shown as "days-hours:minutes". One of the daughter cells dies at time point 3-06:42 (red asterisk), whereas the second cell (dashed box) undergoes progressive morphological changes finally acquiring a neuronal morphology and TuJ1 expression (not shown) after 5 days.

5.3. Recruitment of local or SEZ astroglia following injury of the postnatal cortex?

While the present data argue for a high degree of plasticity of early postnatal astroglia in culture, astroglia fate-mapping studies from the Vaccarino lab suggest that to some low degree (<1%) astroglia can spontaneously generate neurons even in the postnatal cortex in vivo (Ganat et al., 2006). Moreover, following hypoxic insult, which causes a marked reduction in cortical thickness and neuron number at early postnatal stages, a much more substantial incorporation of new neurons can be observed (Fagel et al., 2009, 2006) which may eventually result in total recovery of cortical size and cell number. However, it is currently not known whether these newly incorporated neurons are derived from astroglia endogenous to the injured cortex or are rather from astroglial cells residing within the emerging adult neurogenic SEZ, which responds to hypoxic injury with a marked increase in proliferation (Fagel et al., 2009, 2006; Sundholm-Peters et al., 2005; Szele and Chesselet, 1996). The fact that many of the newly generated neurons were found to express Tbr1 (Fagel et al., 2006), as mentioned above a hallmark of glutamatergic

neurogenesis, may on first sight argue for local generation, as postnatally generated neurons in the SEZ comprise largely GABAergic neurons. However, a recent study has demonstrated that the adult SEZ also contains a pool of glutamatergic progenitors which can be recruited from the SEZ upon injury (Brill et al., 2009).

6. Re-acquisition of a stem or progenitor-like state by mature glia following injury

6.1. Reactivation of astroglia following injury of the cerebral cortex

Thus, several lines of evidence suggest that the abysm between RGCs and early postnatal astroglia is not yet unbridgeable, but becomes wider and wider with development. Upon reaching adulthood at the latest, however, astrocytes resident to the cortical parenchyma seem to have lost all neurogenic potential and accordingly the adult cortex is devoid of newly generated neurons. Recently, Buffo and et al. re-assessed the question whether astroglia proliferates in

the mature cerebral cortex by genetic fate-mapping, using a mouse line expressing a tamoxifen-inducible Cre recombinase driven by the astrocyte-specific GLAST promoter crossed to a reporter line (Buffo et al., 2008). Notably, at that stage astroglia was found to become largely quiescent and hardly any genetically fate-mapped astroglia incorporated the thymidine analogue BrdU. Similar data were also obtained by doublelabelling immunohistochemical makers for astrocytes with BrdU (Buffo et al., 2008) (C. Simon and M. Götz, unpublished observation). This situation changes dramatically following injury. When the cerebral cortex was locally injured by a stab wound, genetically fate-mapped astroglia was found to resume proliferation along side with the up-regulation of GFAP and other classical markers of reactive glia. Despite the observation that some injury paradigms appear to be inductive for low degree neurogenesis in the cerebral cortex (Magavi et al., 2000), no new neurons could be found in the stab wound model (Buffo et al., 2008, 2005). In fact, virtually all fatemapped astroglia were found to give rise to new astrocytes (Buffo et al., 2008). Does this mean that these proliferating astroglia are intrinsically restricted to the astroglial lineage? In fact, in vivo the answer to this question seems to be yes. Surprisingly, however, when these astroglial cells were placed in vitro, these, but not their counterparts of the intact cortical hemisphere, could give rise to neurospheres characterized by the ability to generate secondary and tertiary neurospheres, suggestive of self-renewal capacity, and by multipotency, i.e. the ability to generate not only new astroglia, but also oligodendrocytes and neurons (Buffo et al., 2008). Intriguingly, ectopic grafting of committed neuronal precursors derived from the SEZ into a non-neurogenic milieu such as the adult striatum causes their glial conversion (Seidenfaden et al., 2006) suggesting that the lack of neurogenesis from reactivated astroglia in vivo may not be due to cell-intrinsic shortcomings of these cells compared to astro/radial glial stem cells of the SEZ, but their exposure to a highly nonneurogenic microenvironment.

Notably, data from our laboratory show that these neurosphere cells can be efficiently reprogrammed in vitro to generate glutamatergic and GABAergic neurons by forced expression of Neurog2 and Dlx2, respectively (Heinrich et al., unpublished observation). Can these cells also be reprogrammed in vivo? The answer to this question is a partial yes. Retrovirally-mediated forced expression of Pax6 was found to induce doublecortin (DCX) in some of the transduced progeny suggesting early stages of neurogenesis, but this population of DCX positive cells eventually disappeared (Buffo et al., 2005). This abortive neurogenesis may have many reasons, such as only partial transdifferentiation leading to failure of functional integration and ultimately apoptosis, therein resembling the high incidence of cell death following recruitment of DCX positive cells from the SEZ into the ischemic striatum (Arvidsson et al., 2002). This data suggests that successful reprogramming of mature astroglia represents only one of the hurdles to be taken if aiming at functional neurological reconstitution of damaged brain circuits. At the same time the microenvironment must be made favourable or at least permissive. Suffice it to say that neurons generated from reprogrammed astroglia must be able to form functional synaptic connections between themselves and the adjacent intact tissue, a process which will require the help of astroglia, well known to play a fundamental role in regulating synapse formation (Christopherson et al., 2005; Slezak and Pfrieger, 2003; Song et al., 2002) and plasticity (Perea and Araque, 2007).

Studies performed in our laboratory clearly indicate that while some early stages of neurogenesis can be indeed attained by forced expression of single neurogenic transcription factors, the response falls short in comparison to the effect observed in early postnatal astroglia. Yet, hope for improvement originates from work in adult pancreas. The Melton laboratory could show that while forced expression of single transcription factors failed to induce an effective reprogramming of exocrine cells into insulin secreting beta cells, a quite substantial degree of transdifferentiation was observed in the adult pancreas in vivo following expression of three defined factors involved in distinct stages of beta cell specification and differentiation (Zhou et al., 2008). It is thus conceivable that joint expression of several neurogenic transcription factors may be more effective in achieving stable lineage transgression towards neuronal identity and thereby also increase the chances for functional integration which in case of incomplete neuronal reprogramming will remain highly compromised.

Although the neurosphere-forming capacity seems to originate from reactive astroglia, we do not know whether all types of parenchymal astroglia de-differentiate to the same degree as to re-acquire NSC like properties. Intriguingly, however there are some hints to which molecular pathways may account for this remarkable response. As mentioned above, the morphogen Shh can induce GFAP-positive cells from the adult cerebral cortex to generate neurospheres (Jiao and Chen, 2008). A recent study has now revealed that Shh expression is indeed induced within cortical astroglia following a local injury {Amankulor, 2009 #267}. Surprisingly, the induction of Shh expression was found to depend on astroglia-macrophage interactions. There is an obvious need to gain more insights into the molecular processes underlying reactive gliosis. Indeed, recent progress unravelled a key role of astroglia-basement membrane contact mediated by integrins in maintaining astroglia in a non-reactive state (Robel et al., 2009). Of greatest interest in regard to the theme of astroglial de-differentiation into stem cells, loss of β 1integrin-mediated signalling leads to the activation of all hallmarks of reactive astrocytes except proliferation and stem cell de-differentiation, thereby highlighting that specific aspects of reactive gliosis are elicited by distinct signalling pathways.

6.2. Reactivation of Müller glia following retinal injury

Intriguingly, a similar response to injury can be observed in the retina. In teleost fish, substantial regeneration of the neural retina can be spontaneously achieved from Müller glia which may be a pendant to astroglia within other central nervous tissues (Lamba et al., 2008a). Indeed, Müller glia in the teleost retina have a complex response to local injury that includes some features of reactive gliosis (up-regulation of glial fibrillary acidic protein, GFAP, and re-entry into the cell cycle) along with characteristics associated with radial glia (expression of BLBP) and re-acquisition of molecular characteristics of multipotent retinal progenitors, such as activation of Notch-Delta signalling and expression of Pax6 (Raymond et al., 2006). The regenerative capacity of the neural retina appears to be progressively lost during evolution. In birds, following acute damage to the neural retina Müller glia still re-enter the cell cycle with many expressing transcription factors characteristic of embryonic retinal progenitors such as Pax6 (Fischer and Reh, 2001). Notably, some of these progenitors then give rise to new neurons, while others produce new Müller glia or remain progenitors. Thus, despite some regenerative capacity, the response remains limited. In mammals such as mice, the regenerative response to neurotoxic injury resulting in the selective death of amacrine and retinal ganglion cells is even more restricted as Müller glia does not spontaneously enter the cell cycle (Karl et al., 2008). However, following intraocular injection of growth factors such as EGF, FGF2 or FGF2 in combination with IGF, it is possible to stimulate Müller glia proliferation and de-differentiation, which now do express Pax6 (Karl et al., 2008). Of note, this up-regulation of Pax6 is in strong contrast to what can be observed in reactive astroglia in the cerebral cortex where instead the basic helix loop helix transcription factor Olig2 is up-regulated, persistent expression of which is antineurogenic in the telencephalon (Buffo et al., 2005; Colak et al., 2008; Hack et al., 2005). Turning back to the retina, consistent with the induction of Pax6, when stimulated with growth factors Müller glia even generate some amacrine neurons. Again, many of these newly generated neurons die (Karl et al., 2008). One may speculate that this could be in part due to the fact that no new retinal ganglion cells are regenerated and thus functional integration of newly generated amacrine neurons remains incomplete eventually leading to their apoptosis.

Using the retinal injury model the Reh laboratory has provided some intriguing evidence to what actually happens when reactive glia de-differentiate. They found that Müller glia start to re-express a component of the SWI/SNF core complex called BAF60c (Lamba et al., 2008b). The SWI/SNF complex is an ATP dependent chromatin remodelling complex that undergoes changes in its composition in parallel to the differentiation from neural stem cell to neuron or glia. Notably, BAF60c is normally expressed in retinal progenitors and re-expression of this factor by Müller glia following injury combined with growth factor treatment suggests that the target specificity and hence remodelling activity SWI/SNF complex becomes progenitor-alike. One particular interesting aspect of BAF60c is its role in stabilizing the interaction between activated Notch and its DNA-binding partner RBP-J (Takeuchi et al., 2007). Thus, re-expression of BAF60c may render Müller glial cells more susceptible to Notch signalling. Intriguingly, in birds Notch signalling seems to play a dual role, namely it contributes to the de-differentiation process, but when maintained at high levels counteracts neurogenesis (Hayes et al., 2007) not unlike to what has been stated above about the role of Notch signalling in RGCs of the developing cerebral cortex.

Taken together, reactive gliosis in the cerebral cortex and the retina results in the re-acquisition of stem cell or progenitor-like properties, but in order to promote a regeneration eventually leading to the reconstitution of neurological function, the details of the reactive response need to be much better understood.

7. The concept of a transcriptional network underlying neural stemcellness

The fact that neural stem and progenitor cells often come in the disguise of glia leads us to one particularly perplexing question: what are the common denominators of the transcriptional core network regulating neural stem cell and (astro)-glial identity? Under transcriptional core network we understand a set of transcription factors which mutually regulate each other, thereby stabilizing the differentiation status of a cell. The concept of a transcriptional core network has been extremely fruitful in the case of embryonic stem cells which eventually led to the discovery of a defined set of transcription factors that allow for the reprogramming of somatic cells (Jaenisch and Young, 2008; Takahashi and Yamanaka, 2006). Forced expression of Oct4, Sox2, Klf4 and c-myc eventually superimposes onto fibroblasts a transcriptional circuitry such that these stably acquire the status of pluripotency. Is there a similar transcriptional circuitry underlying the differentiation state of a "neural stem cell"? The observation that neural stem cells are faterestricted in a region-specific manner, at least with respect to neuronal subtype specification (Merkle et al., 2007), may suggest that there is indeed not a single transcriptional core network conveying the status of a neural stem cell, but potentially more than one. Yet, SEZ stem cells can be easily forced to acquire other neuronal subtype identity following forced expression of appropriate neurogenic fate determinants (Berninger et al., 2007b; Brill et al., 2008; Hack et al., 2005) indicating that they possess the competence to correctly interpret these transcriptional cues. These may argue in favour of a neural stem cell core network that is not region-specific, but universal. Furthermore, some transcription factors such as Sox2 (Cavallaro et al., 2008; Favaro et al., 2009; Graham et al., 2003; Kessaris et al., 2006; Suh et al., 2007; Taranova et al., 2006), Tlx (Liu et al., 2008; Roy et al., 2004; Shi et al., 2004) and Bmi-1 (Fasano et al., 2009; Molofsky et al., 2005, 2003; Moon et al., 2008; Zencak et al., 2005) appear to play a critical role in the maintenance and self-renewal of NSCs of different regions and developmental stages, suggesting that these might be either constituents of a NSC transcriptional network or at least closely related to it. A functional proof for the existence of a NSC transcriptional core network would be provided if forced expression of transcription factors defining this network could superimpose a NSC fate on parenchymal astroglia or even other types of somatic cells that are ontogenetically more distant than astrocytes from NSCs.

Two aspects merit further consideration: first the surprising closeness of neural to pluripotent stem cells. While fibroblast require several transcription factors for reprogramming into induced pluripotent stem cells, sole expression of Oct4 is sufficient to reprogram mouse adult as well as human foetal neural stem cells into pluripotent cells (Kim et al., in press, 2009). These data suggest that the stable transition between a neural and a pluripotent stem cell core networks may be rather simple, and reflect the other side of the coin that neural determination often appears a default pathway of embryonic stem cell differentiation (Tropepe et al., 2001). Such facile transition may reflect a high degree of commonalities between these core networks, as exemplified by the common expression of Sox2.

The second point deals with the question posed at the beginning of the paragraph: How can a NSC core network coexist with a hypothetical transcriptional core network conveying "glianess" within a single cell? Obviously, a NSC transcriptional network is not active in all astroglia, as classical parenchyma astroglia are not functioning as NSCs. However, some candidate constituents like Sox2 are expressed not only in NSCs, but also in parenchymal astroglia (Komitova and Eriksson, 2004). Other candidate genes such as Tlx seem to be restricted to stem cell astroglia (Liu et al., 2008), and loss of Bmi-1 even increases the genesis of non-stem cell astroglia at the expense of NSCs (Zencak et al., 2005), while forced expression of Bmi-1 induces stem cell characteristics in cultured astroglia (Moon et al., 2008). Thus, there seems to be a clear partitioning between stem cell and parenchymal astroglia. Yet, as the studies in the injured cerebral cortex or retina have shown, parenchymal astroglia have the capacity to regain NSCs characteristics (Buffo et al., 2008; Karl et al., 2008). Possibly, this capacity is due to a partial overlap of the two otherwise distinct transcriptional circuits, as represented by the common expression of Sox2, which may enable astroglia in some circumstances to switch from one to the other mode of being, i.e. being stem or parenchymal glia.

Acknowledgment

The authors would like to thank Dr. Christophe Heinrich for discussion and comments on the manuscript. Work by the authors was supported by the DFG, the BMBF and the Bavarian State Ministry of Sciences, Research and the Arts (for NeuroCell).

$\mathsf{R} \to \mathsf{F} \to \mathsf{R} \to$

- Ahn, S., Joyner, A.L., 2005. In vivo analysis of quiescent adult neural stem cells responding to Sonic hedgehog. Nature 437, 894–897.
- Alexson, T.O., Hitoshi, S., Coles, B.L., Bernstein, A., van der Kooy, D., 2006. Notch signaling is required to maintain all neural stem cell populations—irrespective of spatial or temporal niche. Dev. Neurosci. 28, 34–48.
- Alvarez-Buylla, A., Garcia-Verdugo, J.M., Tramontin, A.D., 2001. A unified hypothesis on the lineage of neural stem cells. Nat. Rev. Neurosci. 2, 287–293.
- Alves, J.A., Barone, P., Engelender, S., Froes, M.M., Menezes, J.R., 2002. Initial stages of radial glia astrocytic transformation in the early postnatal anterior subventricular zone. J. Neurobiol. 52, 251–265.
- Androutsellis-Theotokis, A., Leker, R.R., Soldner, F., Hoeppner, D.J., Ravin, R., Poser, S.W., Rueger, M.A., Bae, S.K., Kittappa, R., McKay, R.D., 2006. Notch signalling regulates stem cell numbers in vitro and in vivo. Nature 442, 823–826.
- Anthony, T.E., Heintz, N., 2008. Genetic lineage tracing defines distinct neurogenic and gliogenic stages of ventral telencephalic radial glial development. Neural Dev. 3, 30.
- Anthony, T.E., Klein, C., Fishell, G., Heintz, N., 2004. Radial glia serve as neuronal progenitors in all regions of the central nervous system. Neuron 41, 881–890.

- Arumugam, T.V., Chan, S.L., Jo, D.G., Yilmaz, G., Tang, S.C., Cheng, A., Gleichmann, M., Okun, E., Dixit, V.D., Chigurupati, S., Mughal, M.R., Ouyang, X., Miele, L., Magnus, T., Poosala, S., Granger, D.N., Mattson, M.P., 2006. Gamma secretase-mediated Notch signaling worsens brain damage and functional outcome in ischemic stroke. Nat. Med. 12, 621–623.
- Arvidsson, A., Collin, T., Kirik, D., Kokaia, Z., Lindvall, O., 2002. Neuronal replacement from endogenous precursors in the adult brain after stroke. Nat. Med. 8, 963–970.
- Balordi, F., Fishell, G., 2007. Hedgehog signaling in the subventricular zone is required for both the maintenance of stem cells and the migration of newborn neurons. J. Neurosci. 27, 5936–5947.
- Berninger, B., Costa, M.R., Koch, U., Schroeder, T., Sutor, B., Grothe, B., Götz, M., 2007a. Functional properties of neurons derived from in vitro reprogrammed postnatal astroglia. J. Neurosci. 27, 8654–8664.
- Berninger, B., Guillemot, F., Götz, M., 2007b. Directing neurotransmitter identity of neurones derived from expanded adult neural stem cells. Eur. J. Neurosci. 25, 2581–2590.
- Bertrand, N., Castro, D.S., Guillemot, F., 2002. Proneural genes and the specification of neural cell types. Nat. Rev. Neurosci. 3, 517–530.
- Bordey, A., Sontheimer, H., 1997. Postnatal development of ionic currents in rat hippocampal astrocytes in situ. J. Neurophysiol. 78, 461–477.
- Brill, M.S., Ninkovic, J., Winpenny, E., Hodge, R.D., Ozen, I., Yang, R., Lepier, A., Gascon, S., Erdelyi, F., Szabo, G., Parras, C., Guillemot, F., Frotscher, M., Berninger, B., Hevner, R.F., Raineteau, O., Gotz, M., 2009. Adult generation of glutamatergic olfactory bulb interneurons. Nat. Neurosci. 12, 1524–1533.
- Brill, M.S., Snapyan, M., Wohlfrom, H., Ninkovic, J., Jawerka, M., Mastick, G.S., Ashery-Padan, R., Saghatelyan, A., Berninger, B., Götz, M., 2008. A dlx2- and pax6-dependent transcriptional code for periglomerular neuron specification in the adult olfactory bulb. J. Neurosci. 28, 6439–6452.
- Buffo, A., Rite, I., Tripathi, P., Lepier, A., Colak, D., Horn, A.P., Mori, T., Gotz, M., 2008. Origin and progeny of reactive gliosis: a source of multipotent cells in the injured brain. Proc. Natl. Acad. Sci. U. S. A. 105, 3581–3586.
- Buffo, A., Vosko, M.R., Erturk, D., Hamann, G.F., Jucker, M., Rowitch, D., Gotz, M., 2005. Expression pattern of the transcription factor Olig2 in response to brain injuries: implications for neuronal repair. Proc. Natl. Acad. Sci. U. S. A. 102, 18183–18188.
- Carlen, M., Meletis, K., Goritz, C., Darsalia, V., Evergren, E., Tanigaki, K., Amendola, M., Barnabe-Heider, F., Yeung, M.S., Naldini, L., Honjo, T., Kokaia, Z., Shupliakov, O., Cassidy, R.M., Lindvall, O., Frisen, J., 2009. Forebrain ependymal cells are Notch-dependent and generate neuroblasts and astrocytes after stroke. Nat. Neurosci. 12, 259–267.
- Cavallaro, M., Mariani, J., Lancini, C., Latorre, E., Caccia, R., Gullo, F., Valotta, M., DeBiasi, S., Spinardi, L., Ronchi, A., Wanke, E., Brunelli, S., Favaro, R., Ottolenghi, S., Nicolis, S.K., 2008.
 Impaired generation of mature neurons by neural stem cells from hypomorphic Sox2 mutants. Development 135, 541–557.
- Chenn, A., Walsh, C.A., 2002. Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. Science 297, 365–369.
- Chojnacki, A.K., Mak, G.K., Weiss, S., 2009. Identity crisis for adult periventricular neural stem cells: subventricular zone astrocytes, ependymal cells or both? Nat. Rev. Neurosci. 10, 153–163.
- Christopherson, K.S., Ullian, E.M., Stokes, C.C., Mullowney, C.E., Hell, J.W., Agah, A., Lawler, J., Mosher, D.F., Bornstein, P., Barres, B.A., 2005. Thrombospondins are astrocyte-secreted proteins that promote CNS synaptogenesis. Cell 120, 421–433.
- Colak, D., Mori, T., Brill, M.S., Pfeifer, A., Falk, S., Deng, C., Monteiro, R., Mummery, C., Sommer, L., Gotz, M., 2008. Adult neurogenesis requires Smad4-mediated bone morphogenic protein signaling in stem cells. J. Neurosci. 28, 434–446.

- Costa, M.R., Bucholz, O., Schroeder, T., Gotz, M., 2009. Late origin of glia-restricted progenitors in the developing mouse cerebral cortex. Cereb. Cortex 19 (Suppl 1), i135–i143.
- Dang, L., Yoon, K., Wang, M., Gaiano, N., 2006. Notch3 signaling promotes radial glial/progenitor character in the mammalian telencephalon. Dev. Neurosci. 28, 58–69.
- Dessaud, E., McMahon, A.P., Briscoe, J., 2008. Pattern formation in the vertebrate neural tube: a sonic hedgehog morphogen-regulated transcriptional network. Development 135, 2489–2503.
- Fagel, D.M., Ganat, Y., Cheng, E., Silbereis, J., Ohkubo, Y., Ment, L.R., Vaccarino, F.M., 2009. Fgfr1 is required for cortical regeneration and repair after perinatal hypoxia. J. Neurosci. 29, 1202–1211.
- Fagel, D.M., Ganat, Y., Silbereis, J., Ebbitt, T., Stewart, W., Zhang, H., Ment, L.R., Vaccarino, F.M., 2006. Cortical neurogenesis
- enhanced by chronic perinatal hypoxia. Exp. Neurol. 199, 77–91. Fasano, C.A., Phoenix, T.N., Kokovay, E., Lowry, N., Elkabetz, Y., Dimos, J.T., Lemischka, I.R., Studer, L., Temple, S., 2009. Bmi-1
- cooperates with Foxg1 to maintain neural stem cell self-renewal in the forebrain. Genes Dev. 23, 561–574. Favaro, R., Valotta, M., Ferri, A.L., Latorre, E., Mariani, J., Giachino,
- C., Lancini, C., Tosetti, V., Ottolenghi, S., Taylor, V., Nicolis, S.K., 2009. Hippocampal development and neural stem cell maintenance require Sox2-dependent regulation of Shh. Nat. Neurosci. 12, 1248–1256.
- Fischer, A.J., Reh, T.A., 2001. Muller glia are a potential source of neural regeneration in the postnatal chicken retina. Nat. Neurosci. 4, 247–252.
- Gaiano, N., Nye, J.S., Fishell, G., 2000. Radial glial identity is promoted by Notch1 signaling in the murine forebrain. Neuron 26, 395–404.
- Ganat, Y.M., Silbereis, J., Cave, C., Ngu, H., Anderson, G.M., Ohkubo, Y., Ment, L.R., Vaccarino, F.M., 2006. Early postnatal astroglial cells produce multilineage precursors and neural stem cells in vivo. J. Neurosci. 26, 8609–8621.
- Ge, W., Martinowich, K., Wu, X., He, F., Miyamoto, A., Fan, G., Weinmaster, G., Sun, Y.E., 2002. Notch signaling promotes astrogliogenesis via direct CSL-mediated glial gene activation. J. Neurosci. Res. 69, 848–860.
- Gotz, M., Stoykova, A., Gruss, P., 1998. Pax6 controls radial glia differentiation in the cerebral cortex. Neuron 21, 1031–1044.
- Graham, V., Khudyakov, J., Ellis, P., Pevny, L., 2003. SOX2 functions to maintain neural progenitor identity. Neuron 39, 749–765.
- Hack, M.A., Saghatelyan, A., de Chevigny, A., Pfeifer, A., Ashery-Padan, R., Lledo, P.M., Gotz, M., 2005. Neuronal fate determinants of adult olfactory bulb neurogenesis. Nat. Neurosci. 8, 865–872.
- Han, Y.G., Spassky, N., Romaguera-Ros, M., Garcia-Verdugo, J.M., Aguilar, A., Schneider-Maunoury, S., Alvarez-Buylla, A., 2008. Hedgehog signaling and primary cilia are required for the formation of adult neural stem cells. Nat. Neurosci. 11, 277–284.
- Hartfuss, E., Galli, R., Heins, N., Gotz, M., 2001. Characterization of CNS precursor subtypes and radial glia. Dev. Biol. 229, 15–30.
- Hatakeyama, J., Bessho, Y., Katoh, K., Ookawara, S., Fujioka, M., Guillemot, F., Kageyama, R., 2004. Hes genes regulate size, shape and histogenesis of the nervous system by control of the timing of neural stem cell differentiation. Development 131, 5539–5550.
- Hayes, S., Nelson, B.R., Buckingham, B., Reh, T.A., 2007. Notch signaling regulates regeneration in the avian retina. Dev. Biol. 312, 300–311.
- Heins, N., Malatesta, P., Cecconi, F., Nakafuku, M., Tucker, K.L., Hack, M.A., Chapouton, P., Barde, Y.A., Gotz, M., 2002. Glial cells generate neurons: the role of the transcription factor Pax6. Nat. Neurosci. 5, 308–315.
- Heng, J.I., Nguyen, L., Castro, D.S., Zimmer, C., Wildner, H., Armant, O., Skowronska-Krawczyk, D., Bedogni, F., Matter, J.M., Hevner, R., Guillemot, F., 2008. Neurogenin 2 controls cortical neuron migration through regulation of Rnd2. Nature 455, 114–118.

- Hermanson, O., Jepsen, K., Rosenfeld, M.G., 2002. N-CoR controls differentiation of neural stem cells into astrocytes. Nature 419, 934–939.
- Hevner, R.F., Hodge, R.D., Daza, R.A., Englund, C., 2006. Transcription factors in glutamatergic neurogenesis: conserved programs in neocortex, cerebellum, and adult hippocampus. Neurosci. Res. 55, 223–233.
- Hochstim, C., Deneen, B., Lukaszewicz, A., Zhou, Q., Anderson, D.J., 2008. Identification of positionally distinct astrocyte subtypes whose identifies are specified by a homeodomain code. Cell 133, 510–522.
- Ishibashi, M., Moriyoshi, K., Sasai, Y., Shiota, K., Nakanishi, S., Kageyama, R., 1994. Persistent expression of helix–loop–helix factor HES-1 prevents mammalian neural differentiation in the central nervous system. EMBO J. 13, 1799–1805.
- Jaenisch, R., Young, R., 2008. Stem cells, the molecular circuitry of pluripotency and nuclear reprogramming. Cell 132, 567–582.
- Jiao, J., Chen, D.F., 2008. Induction of neurogenesis in non-conventional neurogenic regions of the adult CNS by niche astrocyte-produced signals. Stem Cells 26, 1221–1230.
- Karl, M.O., Hayes, S., Nelson, B.R., Tan, K., Buckingham, B., Reh, T.A., 2008. Stimulation of neural regeneration in the mouse retina. Proc. Natl. Acad. Sci. U. S. A. 105, 19508–19513.
- Kessaris, N., Fogarty, M., Iannarelli, P., Grist, M., Wegner, M., Richardson, W.D., 2006. Competing waves of oligodendrocytes in the forebrain and postnatal elimination of an embryonic lineage. Nat. Neurosci. 9, 173–179.
- Kim J.B., Greber B., Araúzo-Bravo M.J., Meyer J., Park K.I., Zaehres H., Schöler H.R., in press. Direct reprogramming of human neural stem cells by Oct4, Nature.
- Kim, J.B., Sebastiano, V., Wu, G., Arauzo-Bravo, M.J., Sasse, P., Gentile, L., Ko, K., Ruau, D., Ehrich, M., van den Boom, D., Meyer, J., Hubner, K., Bernemann, C., Ortmeier, C., Zenke, M., Fleischmann, B.K., Zaehres, H., Scholer, H.R., 2009. Oct4-induced pluripotency in adult neural stem cells. Cell 136, 411–419.
- Kohwi, M., Petryniak, M.A., Long, J.E., Ekker, M., Obata, K., Yanagawa, Y., Rubenstein, J.L., Alvarez-Buylla, A., 2007. A subpopulation of olfactory bulb GABAergic interneurons is derived from Emx1- and Dlx5/6-expressing progenitors. J. Neurosci. 27, 6878–6891.
- Komada, M., Saitsu, H., Kinboshi, M., Miura, T., Shiota, K., Ishibashi, M., 2008. Hedgehog signaling is involved in development of the neocortex. Development 135, 2717–2727.
- Komitova, M., Eriksson, P.S., 2004. Sox-2 is expressed by neural progenitors and astroglia in the adult rat brain. Neurosci. Lett. 369, 24–27.
- Kriegstein, A., Alvarez-Buylla, A., 2009. The glial nature of embryonic and adult neural stem cells. Annu. Rev. Neurosci. 32, 149–184.
- Kuwabara, T., Hsieh, J., Muotri, A., Yeo, G., Warashina, M., Lie, D.C., Moore, L., Nakashima, K., Asashima, M., Gage, F.H., 2009.
 Wnt-mediated activation of NeuroD1 and retro-elements during adult neurogenesis. Nat. Neurosci. 12, 1097–1105.
- Lai, K., Kaspar, B.K., Gage, F.H., Schaffer, D.V., 2003. Sonic hedgehog regulates adult neural progenitor proliferation in vitro and in vivo. Nat. Neurosci. 6, 21–27.
- Lamba, D., Karl, M., Reh, T., 2008a. Neural regeneration and cell replacement: a view from the eye. Cell Stem Cell 2, 538–549.
- Lamba, D.A., Hayes, S., Karl, M.O., Reh, T., 2008b. Baf60c is a component of the neural progenitor-specific BAF complex in developing retina. Dev. Dyn. 237, 3016–3023.
- Laywell, E.D., Rakic, P., Kukekov, V.G., Holland, E.C., Steindler, D.A., 2000. Identification of a multipotent astrocytic stem cell in the immature and adult mouse brain. Proc. Natl. Acad. Sci. U. S. A. 97, 13883–13888.
- Li, W., Cogswell, C.A., LoTurco, J.J., 1998. Neuronal differentiation of precursors in the neocortical ventricular zone is triggered by BMP. J. Neurosci. 18, 8853–8862.

Lie, D.C., Colamarino, S.A., Song, H.J., Desire, L., Mira, H., Consiglio, A., Lein, E.S., Jessberger, S., Lansford, H., Dearie, A.R., Gage, F.H., 2005. Wnt signalling regulates adult hippocampal neurogenesis. Nature 437, 1370–1375.

Liu, H.K., Belz, T., Bock, D., Takacs, A., Wu, H., Lichter, P., Chai, M., Schutz, G., 2008. The nuclear receptor tailless is required for neurogenesis in the adult subventricular zone. Genes Dev. 22, 2473–2478.

LoTurco, J.J., Bai, J., 2006. The multipolar stage and disruptions in neuronal migration. Trends Neurosci. 29, 407–413.

Magavi, S.S., Leavitt, B.R., Macklis, J.D., 2000. Induction of neurogenesis in the neocortex of adult mice. Nature 405, 951–955.

Malatesta, P., Hack, M.A., Hartfuss, E., Kettenmann, H., Klinkert, W., Kirchhoff, F., Gotz, M., 2003. Neuronal or glial progeny: regional differences in radial glia fate. Neuron 37, 751–764.

Malatesta, P., Hartfuss, E., Gotz, M., 2000. Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. Development 127, 5253–5263.

Merkle, F.T., Mirzadeh, Z., Alvarez-Buylla, A., 2007. Mosaic organization of neural stem cells in the adult brain. Science 317, 381–384.

Merkle, F.T., Tramontin, A.D., Garcia-Verdugo, J.M., Alvarez-Buylla, A., 2004. Radial glia give rise to adult neural stem cells in the subventricular zone. Proc. Natl. Acad. Sci. U. S. A. 101, 17528–17532.

Miyata, T., Kawaguchi, A., Okano, H., Ogawa, M., 2001. Asymmetric inheritance of radial glial fibers by cortical neurons. Neuron 31, 727–741.

Mizutani, K., Yoon, K., Dang, L., Tokunaga, A., Gaiano, N., 2007. Differential Notch signalling distinguishes neural stem cells from intermediate progenitors. Nature 449, 351–355.

Molofsky, A.V., He, S., Bydon, M., Morrison, S.J., Pardal, R., 2005. Bmi-1 promotes neural stem cell self-renewal and neural development but not mouse growth and survival by repressing the p16Ink4a and p19Arf senescence pathways. Genes Dev. 19, 1432–1437.

Molofsky, A.V., Pardal, R., Iwashita, T., Park, I.K., Clarke, M.F., Morrison, S.J., 2003. Bmi-1 dependence distinguishes neural stem cell self-renewal from progenitor proliferation. Nature 425, 962–967.

Moon, J.H., Yoon, B.S., Kim, B., Park, G., Jung, H.Y., Maeng, I., Jun, E.K., Yoo, S.J., Kim, A., Oh, S., Whang, K.Y., Kim, H., Kim, D.W., Kim, K.D., You, S., 2008. Induction of neural stem cell-like cells (NSCLCs) from mouse astrocytes by Bmi1. Biochem. Biophys. Res. Commun. 371, 267–272.

Noctor, S.C., Flint, A.C., Weissman, T.A., Dammerman, R.S., Kriegstein, A.R., 2001. Neurons derived from radial glial cells establish radial units in neocortex. Nature 409, 714–720.

Noctor, S.C., Flint, A.C., Weissman, T.A., Wong, W.S., Clinton, B.K., Kriegstein, A.R., 2002. Dividing precursor cells of the embryonic cortical ventricular zone have morphological and molecular characteristics of radial glia. J. Neurosci. 22, 3161–3173.

Noctor, S.C., Martinez-Cerdeno, V., Ivic, L., Kriegstein, A.R., 2004. Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. Nat. Neurosci. 7, 136–144.

Olsen, M.L., Sontheimer, H., 2008. Functional implications for Kir4.1 channels in glial biology: from K+ buffering to cell differentiation. J. Neurochem. 107, 589–601.

Palma, V., Lim, D.A., Dahmane, N., Sanchez, P., Brionne, T.C., Herzberg, C.D., Gitton, Y., Carleton, A., Alvarez-Buylla, A., Ruiz i Altaba, A., 2005. Sonic hedgehog controls stem cell behavior in the postnatal and adult brain. Development 132, 335–344.

Perea, G., Araque, A., 2007. Astrocytes potentiate transmitter release at single hippocampal synapses. Science 317, 1083–1086.

Petryniak, M.A., Potter, G.B., Rowitch, D.H., Rubenstein, J.L., 2007. Dlx1 and Dlx2 control neuronal versus oligodendroglial cell fate acquisition in the developing forebrain. Neuron 55, 417–433.

- Pinto, L., Gotz, M., 2007. Radial glial cell heterogeneity—the source of diverse progeny in the CNS. Prog. Neurobiol. 83, 2–23.
- Qian, X., Shen, Q., Goderie, S.K., He, W., Capela, A., Davis, A.A., Temple, S., 2000. Timing of CNS cell generation: a programmed sequence of neuron and glial cell production from isolated murine cortical stem cells. Neuron 28, 69–80.
- Rakic, P., 2003. Elusive radial glial cells: historical and evolutionary perspective. Glia 43, 19–32.

Raymond, P.A., Barthel, L.K., Bernardos, R.L., Perkowski, J.J., 2006. Molecular characterisation of retinal stem cells and their niches in adult zebrafish. BMC Dev. Biol. 6, 36.

Robel, S., Mori, T., Zoubaa, S., Schlegel, J., Sirko, S., Faissner, A., Goebbels, S., Dimou, L., Gotz, M., 2009. Conditional deletion of beta1-integrin in astroglia causes partial reactive gliosis. Glia 57, 1630–1647.

Roy, K., Kuznicki, K., Wu, Q., Sun, Z., Bock, D., Schutz, G., Vranich, N., Monaghan, A.P., 2004. The Tlx gene regulates the timing of neurogenesis in the cortex. J. Neurosci. 24, 8333–8345.

Sakurai, K., Osumi, N., 2008. The neurogenesis-controlling factor, Pax6, inhibits proliferation and promotes maturation in murine astrocytes. J. Neurosci. 28, 4604–4612.

Seidenfaden, R., Desoeuvre, A., Bosio, A., Virard, I., Cremer, H., 2006. Glial conversion of SVZ-derived committed neuronal precursors after ectopic grafting into the adult brain. Mol. Cell. Neurosci. 32, 187–198.

Seifert, G., Huttmann, K., Binder, D.K., Hartmann, C., Wyczynski, A., Neusch, C., Steinhauser, C., 2009. Analysis of astroglial K+ channel expression in the developing hippocampus reveals a predominant role of the Kir4.1 subunit. J. Neurosci. 29, 7474–7488.

Shi, Y., Chichung Lie, D., Taupin, P., Nakashima, K., Ray, J., Yu, R.T., Gage, F.H., Evans, R.M., 2004. Expression and function of orphan nuclear receptor TLX in adult neural stem cells. Nature 427, 78–83.

Shimojo, H., Ohtsuka, T., Kageyama, R., 2008. Oscillations in notch signaling regulate maintenance of neural progenitors. Neuron 58, 52–64.

Slezak, M., Pfrieger, F.W., 2003. New roles for astrocytes: regulation of CNS synaptogenesis. Trends Neurosci. 26, 531–535.

Song, H.J., Stevens, C.F., Gage, F.H., 2002. Neural stem cells from adult hippocampus develop essential properties of functional CNS neurons. Nat. Neurosci. 5, 438–445.

Spassky, N., Merkle, F.T., Flames, N., Tramontin, A.D., Garcia-Verdugo, J.M., Alvarez-Buylla, A., 2005. Adult ependymal cells are postmitotic and are derived from radial glial cells during embryogenesis. J. Neurosci. 25, 10–18.

Suh, H., Consiglio, A., Ray, J., Sawai, T., D'Amour, K.A., Gage, F.H., 2007. In vivo fate analysis reveals the multipotent and self-renewal capacities of Sox2+ neural stem cells in the adult hippocampus. Cell Stem Cell 1, 515–528.

Sundholm-Peters, N.L., Yang, H.K., Goings, G.E., Walker, A.S., Szele, F.G., 2005. Subventricular zone neuroblasts emigrate toward cortical lesions. J. Neuropathol. Exp. Neurol. 64, 1089–1100.

Szele, F.G., Chesselet, M.F., 1996. Cortical lesions induce an increase in cell number and PSA-NCAM expression in the subventricular zone of adult rats. J. Comp. Neurol. 368, 439–454.

Takahashi, K., Yamanaka, S., 2006. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. Cell 126, 663–676.

Takeuchi, J.K., Lickert, H., Bisgrove, B.W., Sun, X., Yamamoto, M., Chawengsaksophak, K., Hamada, H., Yost, H.J., Rossant, J., Bruneau, B.G., 2007. Baf60c is a nuclear Notch signaling component required for the establishment of left–right asymmetry. Proc. Natl. Acad. Sci. U. S. A. 104, 846–851.

Tamamaki, N., Nakamura, K., Okamoto, K., Kaneko, T., 2001. Radial glia is a progenitor of neocortical neurons in the developing cerebral cortex. Neurosci. Res. 41, 51–60.

- Taranova, O.V., Magness, S.T., Fagan, B.M., Wu, Y., Surzenko, N., Hutton, S.R., Pevny, L.H., 2006. SOX2 is a dose-dependent regulator of retinal neural progenitor competence. Genes Dev. 20, 1187–1202.
- Tropepe, V., Hitoshi, S., Mak, T.W., Rossant, J., van der Kooy, D., 2001. Direct neural fate specification from embryonic stem cells: a primitive mammalian neural stem cell stage acquired through a default mechanism. Neuron 30, 65–78.
- Voigt, T., 1989. Development of glial cells in the cerebral wall of ferrets: direct tracing of their transformation from radial glia into astrocytes. J. Comp. Neurol. 289, 74–88.
- Wang, D.D., Bordey, A., 2008. The astrocyte odyssey. Prog. Neurobiol. 86, 342–367.
- Woodhead, G.J., Mutch, C.A., Olson, E.C., Chenn, A., 2006. Cell-autonomous beta-catenin signaling regulates cortical precursor proliferation. J. Neurosci. 26, 12620–12630.

- Yoon, K., Nery, S., Rutlin, M.L., Radtke, F., Fishell, G., Gaiano, N., 2004. Fibroblast growth factor receptor signaling promotes radial glial identity and interacts with Notch1 signaling in telencephalic progenitors. J. Neurosci. 24, 9497–9506.
- Young, K.M., Fogarty, M., Kessaris, N., Richardson, W.D., 2007. Subventricular zone stem cells are heterogeneous with respect to their embryonic origins and neurogenic fates in the adult olfactory bulb. J. Neurosci. 27, 8286–8296.
- Zencak, D., Lingbeek, M., Kostic, C., Tekaya, M., Tanger, E., Hornfeld, D., Jaquet, M., Munier, F.L., Schorderet, D.F., van Lohuizen, M., Arsenijevic, Y., 2005. Bmi1 loss produces an increase in astroglial cells and a decrease in neural stem cell population and proliferation. J. Neurosci. 25, 5774–5783.
- Zhou, Q., Brown, J., Kanarek, A., Rajagopal, J., Melton, D.A., 2008. In vivo reprogramming of adult pancreatic exocrine cells to beta-cells. Nature 455, 627–632.